

# NMDAR2A Rabbit mAb

货号: B28680

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	144kDa/165kDa
实测分子量	165kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	Cell junction,Cell membrane,Multi-pass membrane protein,postsynaptic cell membrane,synapse
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NMDAR2A.
序列	LDPPQFVENVTLPENVDFDPYQDPSENFRRKGDSTLPMNRNPLHNEEGLSNNDQYKLYSKHFTLKDKGSPHSETSERYR QNSTHCRSCLSNMPTYSGHFTMRSPFKCDACLRMGNLYDIDEDQMLQETGNPATGEQVYQQDWAQNNALQLQKNKLR ISRQHSYDNIVDKPRELDLSRPSRSISLKDRERLLEGNFYGSLSVPSSKLSGKKSSLPQGLEDSKRKSLLPDHTSDNPFL HSHRDDQRLVIGRCPSDPYKHSLPSQAVNDSY

靶点信息

研究背景	This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is a n N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without mental retardation. Alternative splicing results in multiple transcript variants.
基因ID	2903
基因名	GRIN2A
Swiss	Q12879
别名	EPND;FESD;GluN2A;LKS;NMDAR2A;NR2A;GRIN2A;NMDA 2A

产品验证



实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)