

# **RNA Helicase A Rabbit mAb**

货号: B28706

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	141kDa
实测分子量	141kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse testis
细胞定位	Cytoplasm,Nucleus,nucleolus
纯化	Affinity purification

# 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human RNA Helicase A.
序列	ITGLRAAMEALVVEVTKQPAIISQLDPVNERMLNMIRQISRPSAAGINLMIGSTRYGDGPRPPKMARYDNGSGYRRGGSSY SGGGYGGGYSSGGYGS

### 靶点信息

研究背景	This gene encodes a member of the DEAH-containing family of RNA helicases. The encoded protein is an enzyme that catalyzes the ATP-dependent unwinding of double-stranded RNA and DNA-RNA complexes. This protein localizes to both the nucleus and the cytoplasm and functions as a transcriptional regulator. This protein may also be involved in the expression and nuclear export of retroviral RNAs. Alternate splicing results in multiple transcript variants. Pseudogenes of this gene are found on chromosomes 11 and 13.[ provided by RefSeq, Feb 2010]

基因ID	1660
基因名	DHX9
Swiss	Q08211
别名	DDX9; LKP; NDH2; NDHII; RHA

# 产品验证



Western blot analysis of RNA Helicase A expressed in HeLa,Mouse testis using RNA Helicase A Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn