

# **CRABP2 Rabbit mAb**

货号: B28719

## 产品信息

Human,Mouse,Rat
Rabbit
Monoclonal
WB IHC IF/ICC FC
WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
15kDa
14kDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
Unconjugated
MCF7,SW480,A-431,22Rv1,HT-29,293T,Mouse brain,Mouse uterus
Cytoplasm,Endoplasmic reticulum,Nucleus
Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CRABP2.
序列	VLGVNVMLRKIAVAAASKPAVEIKQEGDTFYIKTSTTVRTTEINFKVGEEFEEQTVDGRPCKSLVKWESENKM

## 靶点信息

研究背景	This gene encodes a member of the retinoic acid (RA, a form of vitamin A) binding protein family and lipo calin/cytosolic fatty-acid binding protein family. The protein is a cytosol-to-nuclear shuttling protein, which facilitates RA binding to its cognate receptor complex and transfer to the nucleus. It is involved in the retinoid signaling pathway, and is associated with increased circulating low-density lipoprotein cholesterol. Alternatively spliced transcript variants encoding the same protein have been found for this gene.
------	--

基因ID	1382
基因名	CRABP2
Swiss	P29373
别名	CRABP2;CRABP-II;RBP6

## 产品验证



Western blot analysis of CRABP2 expressed in Mouse brain,MCF7,SW480 using CRABP2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lan e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn