

SPOP Rabbit mAb

货号: B28755

产品信息

| | |
|-------|---|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 42kDa |
| 实测分子量 | 42kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | PC-3 |
| 细胞定位 | cytoplasm,nuclear speck,nucleoplasm,nucleus |
| 纯化 | Affinity purification |

抗原信息


| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human SPOP. |
| 序列 | VFKEMMCFIYTGKAPNLDKMADDLLAAADKYALERLKVMCEDALCSNLSVENAAEILILADLHSADQLKTQAVDFINYHAS DVLETSGWKSMVVSHPHLVA |

靶点信息

| | |
|------|---|
| 研究背景 | This gene encodes a protein that may modulate the transcriptional repression activities of death-associated protein 6 (DAXX), which interacts with histone deacetylase, core histones, and other histone-associated proteins. In mouse, the encoded protein binds to the putative leucine zipper domain of macroH2A1.2, a variant H2A histone that is enriched on inactivated X chromosomes. The BTB/POZ domain of this protein has been shown in other proteins to mediate transcriptional repression and to interact with components of histone deacetylase co-repressor complexes. Alternative splicing of this gene results in multiple transcript variants encoding the same protein. [provided by RefSeq, Jul 2008] |
|------|---|

| | |
|-------|--------------|
| 基因ID | 8405 |
| 基因名 | SPOP |
| Swiss | O43791 |
| 别名 | BTBD32; TEF2 |

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn