

LI Cadherin Rabbit mAb

货号: B28807

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	120kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,Mouse large intestine
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human LI Cadherin.
序列	MILQAHLHSLCLLMLYLATGYGQEGKFSGPLKPMTFSIYEGQEPSQIIFQFKANPPAVTFELTGETDNIFVIEREGLLYYNRAL DRETRSTHNLQVAALD

靶点信息

研究背景	This gene is a member of the cadherin superfamily, genes encoding calcium-dependent, membrane-asso ciated glycoproteins. The encoded protein is cadherin-like, consisting of an extracellular region, containin g 7 cadherin domains, and a transmembrane region but lacking the conserved cytoplasmic domain. The p rotein is a component of the gastrointestinal tract and pancreatic ducts, acting as an intestinal proton-de pendent peptide transporter in the first step in oral absorption of many medically important peptide-base d drugs. The protein may also play a role in the morphological organization of liver and intestine. Alternati ve splicing results in multiple transcript variants. [provided by RefSeq, Jan 2009]
基因ID	1015
基因名	CDH17
Swiss	Q12864
别名	CDH16; HPT-1; HPT1

产品验证



Western blot analysis of LI Cadherin expressed in HT-29, Mouse large intestine using LI Cadherin Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn