

Complement factor B Rabbit mAb

货号: B28809

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	68kDa/85kDa
实测分子量	100kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat serum
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Complement factor B.
序列	GSWSTLKTQDQKTVRKAECRAIHCPRPDHFENGEYWPRSPYYNVSD EISFHCYDGYTLRGSANRTCQVNGRWSGQTAIC DNGAGYCSNPGIPIGTRKVGSGYRLED SVTYHCSRGLTLRGSQRRTCQEGGSWSGTEPSCQDSFMYDTPQEVAEAF LSS LTETIEGVDAEDGHGPGEQQKRKIVLDPSGSMNIYLVLDGSDSIGASNFTGAKKCLVNLIEKVASYGVKPRYGLVTYATYPK IWVKVSEADSSNADWVTQLNEINYEDHKLKSGTNTKKALQAVYSMMSWPDDVPPEGWNRTRHVIILMTDGLHNMGG DPITVIDEIRDLLYIGKDRKNPRE

靶点信息

研究背景	This gene encodes complement factor B, a component of the alternative pathway of complement activation. Factor B circulates in the blood as a single chain polypeptide. Upon activation of the alternative pathway, it is cleaved by complement factor D yielding the noncatalytic chain Ba and the catalytic subunit Bb. The active subunit Bb is a serine protease which associates with C3b to form the alternative pathway C3 convertase. Bb is involved in the proliferation of preactivated B lymphocytes, while Ba inhibits their proliferation. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. This cluster includes several genes involved in regulation of the immune reaction. Polymorphisms in this gene are associated with a reduced risk of age-related macular degeneration. The polyadenylation site of this gene is 421 bp from the 5' end of the gene for complement component 2.
基因ID	629
基因名	CFB
Swiss	P00751
别名	CFB;AHUS4;ARMD14;BF;BFD;CFAB;CFBD;FB;FBI12;GBG;H2-Bf;PBF2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn