

# 5 Lipoxygenase Rabbit mAb

货号: B28839

## 产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	78kDa
实测分子量	78kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Rat lung
细胞定位	Cytoplasm, Nucleus matrix, Nucleus membrane, Peripheral membrane protein
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human 5 Lipoxygenase.
序列	CYRWITGDVEVVLRDGRAKLARDDQIHILKQHRRKELETRQKQYRWMEWNPGFPLSIDAKCHKDLPRDIQFDSEKGVDF VLNYSKAMENLFINRFMHMFQS

靶点信息

研究背景	This gene encodes a member of the lipoxygenase gene family and plays a dual role in the synthesis of leu kotrienes from arachidonic acid. The encoded protein, which is expressed specifically in bone marrow-der ived cells, catalyzes the conversion of arachidonic acid to 5(S)-hydroperoxy-6-trans-8,11,14-cis-eicosatetr aenoic acid, and further to the allylic epoxide 5(S)-trans-7,9-trans-11,14-cis-eicosatetrenoic acid (leukotri ene A4). Leukotrienes are important mediators of a number of inflammatory and allergic conditions. Muta tions in the promoter region of this gene lead to a diminished response to antileukotriene drugs used in the treatment of asthma and may also be associated with atherosclerosis and several cancers. Alternative ly spliced transcript variants encoding different isoforms have been found for this gene. [provided by Ref Seq, Jan 2012]
基因 <b>ID</b>	240
基因名	ALOX5
Swiss	P09917
别名	5-LO; 5-LOX; 5LPG; LOG5

## 产品验证



Western blot analysis of 5 Lipoxygenase expressed in Raji,Rat lung using 5 Lipoxygenase Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn