

CD134 Rabbit mAb

货号: B28842

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	29kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,BT-474,SW620,MCF7,Mouse thymus,Mouse skeletal muscle,Mouse lung,Mouse intestine,Rat liver
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD134.
序列	LHCVGDTYPSNDRCCHECRPGNGMVSRCRSQNTVCRPCPGFYNDVVSSKPKCTWCNLRGSERKQLCTATQDTV CRCRAGTQPLDSYKPGVDCAPCPPGHFSPGDNQACKPWTNCTLAGKHTLQPASNSSDAICEDRDPPATQPQETQGPPA RPITVQPTEAWPRTSQGPSTRPVEPGGRA

靶点信息

研究背景	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2IL1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation.
------	---

基因ID	7293
基因名	TNFRSF4
Swiss	P43489
别名	TNFRSF4;ACT35;CD134;IMD16;OX40;TXGP1L

产品验证



Western blot analysis of CD134 expressed in HepG2,BT-474,SW620 using CD134 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn