

# TRADD Rabbit mAb

货号: **B28863**

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>ICC:</b> 1:50 - 1:200 <b>IP:</b> 1:20 - 1:50
理论分子量	27kDa/34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Mouse spleen,Mouse kidney
细胞定位	Cytoplasm,Nucleus,cytoskeleton
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TRADD .
序列	QTFLFQGQPVVNRPLSLKDKQTFARISVGLKWRKVGRSLQRGCRALRDPALDSLAYEYEREGLYEQAFQLLRRFVQAEGR RATLQRLVEALEENELTS LAEDLLGLTDPNGGLA

## 靶点信息

研究背景	The protein encoded by this gene is a death domain containing adaptor molecule that interacts with TNF RSF1A/TNFR1 and mediates programmed cell death signaling and NF-kappaB activation. This protein binds adaptor protein TRAF2, reduces the recruitment of inhibitor-of-apoptosis proteins (IAPs) by TRAF2, and thus suppresses TRAF2 mediated apoptosis. This protein can also interact with receptor TNFRSF6/FAS and adaptor protein FADD/MORT1, and is involved in the Fas-induced cell death pathway.
------	---

基因ID	8717
基因名	TRADD
Swiss	Q15628
别名	TRADD; Hs.89862; TNFRSF1A associated via death domain

## 产品验证



Western blot analysis of TRADD expressed in Mouse brain, Mouse spleen, Mouse kidney using TRADD Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)