

Fbx32 Rabbit mAb

货号: B28864

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	30kDa/41kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,A-431,MCF7
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Fbx32.
序列	WQQQLNNIQITRPAFKGLTFTDLPLCLQLNIMQRLSDGRDLVSLGQAAPDLHVLSEDRLLWKKLCQYHFSERQIRKRLILS DKGQLDWKKMYFKLVRCYPRKEQYGDTLQLCKHCHILSWKGTDHPCTANNPESCSVSLSPQDFINLFKF

靶点信息

研究背景	This gene encodes a member of the F-box protein family which is characterized by an approximately 40 a mino acid motif, the F-box. The F-box proteins constitute one of the four subunits of the ubiquitin protein I igase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitinatio n. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine -rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbxs class and contains an F-box domain. This protein is highly expressed during muscle atrophy, whereas mice deficient in this gene were found to be re sistant to atrophy. This protein is thus a potential drug target for the treatment of muscle atrophy. Alterna tive splicing results in multiple transcript variants encoding different isoforms.
基因 ID	114907
基因名	FBXO32
Swiss	Q969P5
别名	FBXO32;Fbx32;MAFbx

产品验证



Western blot analysis of Fbx32 expressed in Jurkat,A-431,MCF7 using Fbx32 Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blockin g buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn