

Histone H3 (acetyl K23) Rabbit mAb

货号: B28938

产品信息

展記 Human,Mouse 和主 Rabbit 克隆性 Monoclonal 類測反应 応用 WB IF/ICC ChIP 推荐浓度 WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 理论分子録 17 kDa 実調分子最 15kDa 形式 Liquid 様存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.		
	反应	Human,Mouse
適用 WB IF/ICC ChIP 推荐浓度 WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 理论分子量 17 kDa 实测分子量 15kDa 形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. 供験 Unconjugated	宿主	Rabbit
	克隆性	Monoclonal
# 接	预测反应	
IF/ICC: 1:50 - 1:200 理论分子量 17 kDa 实测分子量 15kDa 形式 Liquid 保存条件 Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.	应用	WB IF/ICC ChIP
実測分子量15kDa形式Liquid保存条件Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.偶联物Unconjugated	推荐浓度	
形式 Liquid 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. (偶联物 Unconjugated	理论分子量	17 kDa
保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. 【图联物 Unconjugated	实测分子量	15kDa
Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.	形式	Liquid
, , ,	保存条件	
	偶联物	Unconjugated
附性对照 HeLa,NIH/3T3,C6	阳性对照	HeLa,NIH/3T3,C6
细胞定位 extracellular exosome,extracellular region,nucleoplasm,nucleus	细胞定位	extracellular exosome,extracellular region,nucleoplasm,nucleus
绝化 Affinity purification	纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Histone H3 (acetyl K23).
序列	AARKS

靶点信息

研究背景	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
基因ID	8350,3020,8290,440093,126961

基因名	HIST3H3,H3F3C
Swiss	P68431/P84243/Q16695/Q6NXT2/Q71DI3
别名	

产品验证



Western blot analysis of Histone H3 (acetyl K23) expressed in HeLa,NIH/3T3,C6 using Histone H3 (acetyl K23) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:50 00. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn