

# PPP2R5E Rabbit mAb

货号: B28942

### 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	46kDa/54kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,NIH/3T3,mouse thymus,mouse kidney
细胞定位	Cytoplasm
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PPP2R5E.
序列	LLKDVPSSEQPELFLKKLQQCCVIFDFMDTLSDLKMKEYKRSTLNELVDYITISRGCLTEQTYPEVVRMVSCNIFRTLPPSDS NEFDPEEDEPTLEASWPH

靶点信息

研究背景	The protein encoded by this gene belongs to the phosphatase 2A regulatory subunit B family. Protein pho sphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catal ytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. The B regulatory subunit might modulate substrate selectivity and catalytic activity. This gene encodes an ep silon isoform of the regulatory subunit B56 subfamily. Multiple transcript variants encoding several different isoforms have been found for this gene.
基因ID	5529
基因名	PPP2R5E
Swiss	Q16537
别名	PPP2R5E;B56E;B56epsilon

### 产品验证



Western blot analysis of PPP2R5E expressed in HeLa,Jurkat,NIH/3T3 using PPP2R5E Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn