

# TIRAP Rabbit mAb

货号: B28983

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	23kDa/25kDa/27kDa
实测分子量	28kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Cell membrane,Cytoplasm,Membrane
纯化	Affinity purification

## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TIRAP.
序列	MASSTSLPAPGSRPKKPLGKMADWFRQTLLKKPKKRPNPESTSSDASQPTSQDSPLPPSLSVTSPSLPPTHASDSGSS RWSKDYDVCVCHSEEDLVAAQDLVSYLEGSTASLRCFLQLRDAATPGGAIVSELCQALSSSHCRVLLITPGFLQDPWCKYQ MLQALTEAPGAEGCTIPLLSGLSRAAYPPELRFMYYVDGRGPDDGGFRQVKEAVMRYLQTLS


## 靶点信息

研究背景	The innate immune system recognizes microbial pathogens through Toll-like receptors (TLRs), which identify pathogen-associated molecular patterns. Different TLRs recognize different pathogen-associated molecular patterns and all TLRs have a Toll-interleukin 1 receptor (TIR) domain, which is responsible for signal transduction. The protein encoded by this gene is a TIR adaptor protein involved in the TLR4 signaling pathway of the immune system. It activates NF-kappa-B, MAPK1, MAPK3 and JNK, which then results in cytokine secretion and the inflammatory response. Alternative splicing of this gene results in several transcript variants; however, not all variants have been fully described.
------	---

基因ID	114609
基因名	TIRAP
Swiss	P58753
别名	TIRAP;BACTS1;Mal;MyD88-2;wyatt

## 产品验证

Western blot analysis of TIRAP expressed in HUVEC, mouse liver, rat lung using TIRAP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



## 实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)