

IRF7 Rabbit mAb

货号: B29034

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	18kDa/51kDa/54kDa/55kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,NIH/3T3
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IRF7.
序列	GVSSLD

靶点信息

研究背景	IRF7 encodes interferon regulatory factor 7, a member of the interferon regulatory transcription factor (IR F) family. IRF7 has been shown to play a role in the transcriptional activation of virus-inducible cellular ge nes, including interferon beta chain genes. Inducible expression of IRF7 is largely restricted to lymphoid ti ssue. Multiple IRF7 transcript variants have been identified, although the functional consequences of thes e have not yet been established.
------	--

基因ID	3665
基因名	IRF7
Swiss	Q92985
别名	IRF7;IMD39;IRF-7H;IRF7A;IRF7B;IRF7C;IRF7H

产品验证



Western blot analysis of IRF7 expressed in 293T,NIH/3T3 using IRF7 Rabbit mAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn