

Thymidylate Synthase Rabbit mAb

货号: B29037

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	26kDa/31kDa/35kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Molt-4
细胞定位	Cytoplasm,Mitochondrion,Mitochondrion inner membrane,Mitochondrion matrix,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Thymidylate Synthase.
序列	MPVAGSELPRRPLPPAAQERDAEPRPPHGELQYLGQIQHILRCGVRKDDRTGTGTLSVFGMQARYSLRDEFLLTTKRVF WKGVL EELLWFIKGSTNAKELSSKGVKIWDANGSRDFLD SLGFSTREEGDLGPVYGFQWRHFGAEYRDMESDYSGQGV DQLQRVIDTIKTNPD RRIMCAWNPRDLPLMALPPCHALCQFYVVNSELSCQLYQRSGDMGLGVPFNIASYALLTYMIAHI TGLKPGDFIHTLGDAHIYLNHIEPLKIQLQREPRFPKLRILRKVEKIDDFKAEDFQIEGYNPHPTIKMEMAV

靶点信息

研究背景	Thymidylate synthase catalyzes the methylation of deoxyuridylate to deoxythymidylate using 5,10-methylenetetrahydrofolate (methylene-THF) as a cofactor. This function maintains the dTMP (thymidine-5-prime monophosphate) pool critical for DNA replication and repair. The enzyme has been of interest as a target for cancer chemotherapeutic agents. It is considered to be the primary site of action for 5-fluorouracil, 5-fluoro-2-prime-deoxyuridine, and some folate analogs. Expression of this gene and that of a naturally occurring antisense transcript rTSalpha (GeneID:55556) vary inversely when cell-growth progresses from late-log to plateau phase.
基因ID	7298
基因名	TYMS
Swiss	P04818
别名	TYMS;HST422;TMS;TS

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn