

Atg4C Rabbit mAb

货号: B29073

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HL-60
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Atg4C.
序列	MEATGTDEVDKLKTKFISAWNNMKYSWVLKTKTYFSRNSPVLLLGKCYHFKYEDEDKTLPAESGCTIEDHVIAGNVEEFRK DFISRIWLTYREEFPQIEGSALTTDCGWGCTLRTGQMLLAQGLILHFLGRAWTWPDALNIENSDSESWTSHTVKKFTASFE ASLSGEREFKTPTISLKETIGKYSDDHEMRNEVYHRKI

靶点信息

研究背景	Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracell ularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentia tion, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encode d protein is also designated as a member of the C-54 family of cysteine proteases. Alternate transcription al splice variants, encoding the same protein, have been characterized.
基因ID	84938
基因名	ATG4C
Swiss	Q96DT6
别名	ATG4C;APG4-C;APG4C;AUTL1;AUTL3

产品验证



Western blot analysis of Atg4C expressed in Jurkat,HL-60 using Atg4C Rabbit mAb at 1:1000. Secondary a ntibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn