

# **FOXA2 Rabbit mAb**

货号: B29137

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Rat lung
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human FOXA2.
序列	MLGAVKMEGHEPSDWSSYYAEPEGYSSVSNMNAGLGMNGMNTYMSMSAAAMGSGSGNMSAGSMNMSSYVGAGMS PSLAGMSPGAGAMAGMGGSAGAAGVA

## 靶点信息

研究背景	This gene encodes a member of the forkhead class of DNA-binding proteins. These hepatocyte nuclear fa ctors are transcriptional activators for liver-specific genes such as albumin and transthyretin, and they als o interact with chromatin. Similar family members in mice have roles in the regulation of metabolism and in the differentiation of the pancreas and liver. This gene has been linked to sporadic cases of maturity-o nset diabetes of the young. Transcript variants encoding different isoforms have been identified for this g ene. [provided by RefSeq, Oct 2008]
------	--

基因ID	3170
基因名	FOXA2
Swiss	Q9Y261
别名	FOXA2;HNF3B;TCF3B

## 产品验证



Western blot analysis of FOXA2 expressed in HepG2,Rat lung using FOXA2 Rabbit mAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn