

IRF5 Rabbit mAb

货号: B29151

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	17kDa/47kDa/54kDa/56kDa/57kDa
实测分子量	56kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	THP-1, U-937, Rat thymus
细胞定位	Cytoplasm, Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human IRF5.
序列	MNQSIPVAPTPPRRVRLKPWLVAQVNNSCQYPGLQWVNGEKKLFCIPWRHATRGPSQDGDNTIFKAWAKETGKYTEGV DEADPAKWKANLRCALNKSRSDFRLLYDGPRDMPPQPYKIYEVCNSNGPAPTDSQPPEDYSFGAGEEEEEEEELQRMLPSLSTEDVKWPPTLQPPTLRPPTLQPPTLQPPVVLGPPAPDPSPSPL


靶点信息

研究背景	This gene encodes a member of the interferon regulatory factor (IRF) family, a group of transcription factors with diverse roles, including virus-mediated activation of interferon, and modulation of cell growth, differentiation, apoptosis, and immune system activity. Members of the IRF family are characterized by a conserved N-terminal DNA-binding domain containing tryptophan (W) repeats. Alternative promoter use and alternative splicing result in multiple transcript variants, and a 30-nt indel polymorphism (SNP rs60344245) can result in loss of a 10-aa segment.
------	--

基因ID	3663
基因名	IRF5
Swiss	Q13568
别名	IRF5;SLEB10

产品验证

Western blot analysis of IRF5 expressed in THP-1,U-937,Rat thymus using IRF5 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn