

SHIP Rabbit mAb

货号: B29152

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 FC: 1:20 - 1:50 IP: 1:20 - 1:50
理论分子量	109kDa/133kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen
细胞定位	Cytoplasm,Membrane,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human SHIP.
序列	PPCSGSSITEIINPNYMGVGPFGPPMPLHVKQTLSPDQQPTAWSYDQPPKDSPLGPCRGESPPTPPGQPPISPKKFLPSTA NRGLPPRTQESRPSDLGKNA

靶点信息

研究背景	This gene is a member of the inositol polyphosphate-5-phosphatase (INPP5) family and encodes a protein with an N-terminal SH2 domain, an inositol phosphatase domain, and two C-terminal protein interaction d omains. Expression of this protein is restricted to hematopoietic cells where its movement from the cytos ol to the plasma membrane is mediated by tyrosine phosphorylation. At the plasma membrane, the prote in hydrolyzes the 5' phosphate from phosphatidylinositol (3,4,5)-trisphosphate and inositol-1,3,4,5-tetraki sphosphate, thereby affecting multiple signaling pathways. The protein is also partly localized to the nucl eus, where it may be involved in nuclear inositol phosphate signaling processes. Overall, the protein funct ions as a negative regulator of myeloid cell proliferation and survival. Mutations in this gene are associate d with defects and cancers of the immune system. Alternative splicing of this gene results in multiple tran script variants.
基因 ID	3635
基因名	INPP5D
Swiss	Q92835
别名	INPP5D;SHIP;SHIP-1;SHIP1;SIP-145;hp51CN;p150Ship

产品验证



Western blot analysis of SHIP expressed in Mouse spleen using SHIP Rabbit mAb at 1:1000. Secondary ant ibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn