

RNF14 Rabbit mAb

货号: B29168

产品信息

| 反应 | Human,Mouse,Rat |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | |
| 实测分子量 | 54kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | SH-SY5Y,Jurkat,K-562,Mouse placenta,Rat brain |
| 细胞定位 | cytoplasm,cytosol,nucleoplasm,nucleus |
| 纯化 | Affinity purification |

抗原信息

| 抗原信息 | Recombinant fusion protein corresponding to Human RNF14. |
|------|---|
| 序列 | LDLMADVVYCPRPCCQLPVMQEPGCTMGICSSCNFAFCTLCRLTYHGVSPCKVTAEKLMDLRNEYLQADEANKRLLDQRY GKRVIQKALEEMESKEWLEKN |


靶点信息

| TIL | 77 | 115. | 豆 |
|-----|----|------|---|
| | | | |

The protein encoded by this gene contains a RING zinc finger, a motif known to be involved in protein-pro tein interactions. This protein interacts with androgen receptor (AR) and may function as a coactivator th at induces AR target gene expression in prostate. A dominant negative mutant of this gene has been dem onstrated to inhibit the AR-mediated growth of prostate cancer. This protein also interacts with class III u biquitin-conjugating enzymes (E2s) and may act as a ubiquitin-ligase (E3) in the ubiquitination of certain nuclear proteins. Six alternatively spliced transcript variants encoding two distinct isoforms have been re ported. [provided by RefSeq, Jan 2011]

| 基因ID | 9604 |
|-------|-------------------------------------|
| 基因名 | RNF14 |
| Swiss | Q9UBS8 |
| 别名 | RNF14;ARA54;HFB30;HRIHFB2038;TRIAD2 |

产品验证



Western blot analysis of RNF14 expressed in Mouse placenta,Rat brain,SH-SY5Y using RNF14 Rabbit mA b at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn