

HDAC8 Rabbit mAb

货号: B29182

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP FC
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	15kDa/16kDa/17kDa/28kDa/29kDa/31kDa/41kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	C6
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human HDAC8.
序列	MEEPEEPADSGQSLVPVYIYSPEYVSMCDSLAKIPKRASMVHSLIEAYALHKQMRIVKPKVASMEEMATFHTDAYLQHLQK VSQEGDDDHDPDSIEYGLGYDCPATEGIFDYAAAIGGATITAAQCLIDGMCKVAINWSGGWHHAKKDEASGFCYLNDAYL GILRLRRKFERILYVDLDLHHGDGVEDAFSFTSKVMTVSLHKFSPGFFPGTGDVSDVGLGKGRIYSVNVPIQDGIQDEKYY QICESVLKEVYQAFNPKAVVLQLGADTIAGDPMCSFNMTVPVGIGKCLKYILQWQLATLILGGGGYNLANTARCWTYLTGVIL GKTLSEIPDHEFFTAYGPDYVLEITPSCRPDNRNEPHRIQILNYIKGNLKHVV

靶点信息

研究背景	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events . Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class I of the histone deacetylase family. It catalyzes t he deacetylation of lysine residues in the histone N-terminal tails and represses transcription in large mul tiprotein complexes with transcriptional co-repressors. Multiple transcript variants encoding different isof orms have been found for this gene.
基因ID	55869
基因名	HDAC8
Swiss	Q9BY41
别名	HDAC8;CDA07;CDLS5;HD8;HDACL1;MRXS6;RPD3;WTS

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn