

Phospho-Rac1/Cdc42 (Ser71) Rabbit mAb

货号: B29214

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	24kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431
细胞定位	cytoplasm,cytoplasmic ribonucleoprotein granule,cytosol,dendritic spine,extracellular exosome,filopodium,focal adhesion,Golgi transport complex,leading edge membrane,microtubule organizing center,mitotic spindle,phagocytic vesicle,plasma membrane,spindle midzone
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Phospho-Rac1/Cdc42 (Ser71).
序列	PLSYP

靶点信息

研究背景	The protein encoded by this gene is a small GTPase of the Rho-subfamily, which regulates signaling pathways that control diverse cellular functions including cell morphology, migration, endocytosis and cell cycle progression. This protein is highly similar to <i>Saccharomyces cerevisiae</i> Cdc 42, and is able to complement the yeast <i>cdc42-1</i> mutant. The product of oncogene <i>Dbl</i> was reported to specifically catalyze the dissociation of GDP from this protein. This protein could regulate actin polymerization through its direct binding to Neural Wiskott-Aldrich syndrome protein (N-WASP), which subsequently activates Arp2/3 complex. Alternative splicing of this gene results in multiple transcript variants. Pseudogenes of this gene have been identified on chromosomes 3, 4, 5, 7, 8 and 20. [provided by RefSeq, Apr 2013]
基因ID	998,5879
基因名	CDC42,RAC1
Swiss	P60953/P63000
别名	CDC42Hs; G25K; TKS

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn