

TC10 Rabbit mAb

货号: B29250

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	25kDa
实测分子量	25kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,22Rv1,Mouse heart,Rat heart
细胞定位	extracellular exosome,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TC10.
序列	KEEWVPELKEYAPNVPFLLIGTQIDLRDDPKTLARLNDMKEKPICVEQGQKLAKEIGACCYVECSALTQKGLKTVFDEAIIAIL TPKKHTVKKRIGSRCINCCLIT

靶点信息

研究背景	This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho p
	roteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motilit
	y. The encoded protein is an important signalling protein for sarcomere assembly and has been shown to
	play a significant role in the exocytosis of the solute carrier family 2, facilitated glucose transporter mem
	ber 4 and other proteins, possibly acting as the signal that turns on the membrane fusion machinery. Thr
	ee related pseudogene have been identified on chromosomes 2 and 14. [provided by RefSeq, Aug 2011]

基因ID	23433
基因名	RHOQ
Swiss	P17081
别名	ARHQ; HEL-S-42; RASL7A; TC10; TC10A

产品验证



Western blot analysis of TC10 expressed in Rat heart,293T,A-549 using TC10 Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn