

# **IDH1** Rabbit mAb

货号: B29282

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	46kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,C2C12,Mouse liver,Rat liver,Rat ovary,Rat kidney
细胞定位	Cytoplasm,Peroxisome
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human IDH1.
序列	MSKKISGGSVVEMQGDEMTRIIWELIKEKLIFPYVELDLHSYDLGIENRDATNDQVTKDAAEAIKKHNVGVKCATITPDEKR VEEFKLKQMWKSPNGTIRNILGGTVFREAIICKNIPRLVSGWVKPIIIGRHAYGDQYRATDFVVPGPGKVEITYTPSDGTQKV TYLVHNFEEGGGVAMGMYNQDKSIEDFAHSSFQMALSKGWPLYLSTKNTILKKYDGRFKDIFQEIYDKQYKSQFE

靶点信息

研究背景	Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. These e nzymes belong to two distinct subclasses, one of which utilizes NAD(+) as the electron acceptor and the other NADP(+). Five isocitrate dehydrogenases have been reported: three NAD(+)-dependent isocitrate d ehydrogenases, which localize to the mitochondrial matrix, and two NADP(+)-dependent isocitrate dehydr ogenases, one of which is mitochondrial and the other predominantly cytosolic. Each NADP(+)-dependent isozyme is a homodimer. The protein encoded by this gene is the NADP(+)-dependent isocitrate dehydro genase found in the cytoplasm and peroxisomes. It contains the PTS-1 peroxisomal targeting signal sequ ence. The presence of this enzyme in peroxisomes suggests roles in the regeneration of NADPH for intrap eroxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxis omal reactions that consume 2-oxoglutarate, namely the alpha-hydroxylation of phytanic acid. The cytopl asmic enzyme serves a significant role in cytoplasmic NADPH production. Alternatively spliced transcript variants encoding the same protein have been found for this gene.
基因 <b>ID</b>	3417
基因名	IDH1
Swiss	O75874
别名	IDH1;HEL-216;HEL-S-26;IDCD;IDH;IDP;IDPC;PICD

### 产品验证



Western blot analysis of IDH1 expressed in Mouse liver,Rat liver,293T using IDH1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn