

# FABP3 Rabbit mAb

货号: B29298

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	14kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart
细胞定位	Cytoplasm
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human FABP3.
序列	MVDAFLGTWKLVDSKNFDDYMKSLGVGFATRQVASMTKPTTIIEKNGDILTLKTHSTFKNTEISFKLGVEFDETTADDRKV KSIVTLDGGKLVHLQKWDGQETTLVRELIDGKLILTLTHGTAVCTRTYEKEA

## 靶点信息

研究背景	The intracellular fatty acid-binding proteins (FABPs) belongs to a multigene family. FABPs are divided into at least three distinct types, namely the hepatic-, intestinal- and cardiac-type. They form 14-15 kDa prote ins and are thought to participate in the uptake, intracellular metabolism and/or transport of long-chain fa tty acids. They may also be responsible in the modulation of cell growth and proliferation. Fatty acid-bindi ng protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. Thi s gene is a candidate tumor suppressor gene for human breast cancer. Alternative splicing results in mult iple transcript variants.
------	---

基因ID	2170
基因名	FABP3
Swiss	P05413
别名	FABP3;FABP11;H-FABP;M-FABP;MDGI;O-FABP

## 产品验证



Western blot analysis of FABP3 expressed in Mouse heart using FABP3 Rabbit mAb at 1:1000. Secondary a ntibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn