

GAA Rabbit mAb

货号**: B29305**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	120,100,75kDa
实测分子量	120,100,75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,DU145
细胞定位	Lysosome,Lysosome membrane
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human GAA.
序列	QEQCEARGCCYIPAKQGLQGAQMGQPWCFFPPSYPSYKLENLSSSEMGYTATLTRTTPTFFPKDILTLRLDVMMETENRL HFTIKDPANRRYEVPLETPHV

靶点信息

研究背景	This gene encodes lysosomal alpha-glucosidase, which is essential for the degradation of glycogen to glu cose in lysosomes. The encoded preproprotein is proteolytically processed to generate multiple intermedi ate forms and the mature form of the enzyme. Defects in this gene are the cause of glycogen storage dis ease II, also known as Pompe's disease, which is an autosomal recessive disorder with a broad clinical spe ctrum. Alternative splicing results in multiple transcript variants.
基因ID	2548

基因名	GAA
Swiss	P10253
别名	GAA; LYAG; glucosidase alpha; acid

产品验证



Western blot analysis of GAA expressed in 293T,HepG2,DU145 using GAA Rabbit mAb at 1:1000. Secon dary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buf fer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn