

MAP1LC3A Rabbit mAb

货号: B29313

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	16kDa
实测分子量	16kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y
细胞定位	Cytoplasm,Cytoplasmic vesicle,Endomembrane system,Lipid-anchor,autophagosome,autophagosome membrane,cytoskeleton
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MAP1LC3A.
序列	MPSDRPFKQRRSFADRCKEVQQIRDQHPSKIPVIIERYKG EKQLPVLDKTKFLVPDHVNMSELVKIIRRLQLNPTQAFFLLV NQHSMVSVSTPIADIYE

靶点信息

研究背景	MAP1A and MAP1B are microtubule-associated proteins which mediate the physical interactions between microtubules and components of the cytoskeleton. MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. The protein encoded by this gene is one of the light chain subunits and can associate with either MAP1A or MAP1B. Two transcript variants encoding different isoforms have been found for this gene. The expression of variant 1 is suppressed in many tumor cell lines, suggesting that it may be involved in carcinogenesis. [provided by RefSeq, Feb 2012]
基因ID	84557
基因名	MAP1LC3A
Swiss	Q9H492
别名	ATG8E; LC3; LC3A; MAP1ALC3; MAP1BLC3

产品验证



Western blot analysis of MAP1LC3A expressed in SH-SY5Y using MAP1LC3A Rabbit mAb at 1:1000. Second antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn