

Phospho-Histone H2A.X (S139) Rabbit mAb

货号: B29328

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	15kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,NIH/3T3,C2C12,Jurkat,Raji,Rat thymus
细胞定位	Chromosome,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Phospho-Histone H2A.X (S139).
序列	VYLAADVLEYLTAEILELAGNAARDNKKTRIIPRHLQLAIRNDEELNKLGGVTIAQGGVLPNIQAVLLPKKTSATVGPKAPSGG KKATQASQEY

靶点信息

研究背景	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif.
基因ID	3014
基因名	H2AFX
Swiss	P16104
别名	H2A.X;H2A/X;H2AX;Histone H2AX;H2AFX;histone H2AX;

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn