

CTIP1 Rabbit mAb

货号**: B29370**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	120kDa
实测分子量	120kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CTIP1.
序列	GEKPYKCNLCDHACTQASKLKRHMKTHMHKSSPMTVKSDDGLSTASSPEPGTSDLVGSASSALKSVVAKFKSENDPNLI PENGDEEEEEDDEEEEEEEEEEEEEEEEEEEEVDYGFGLSLEAARHHENSSRGAVVGVGDESRALPDVMQGMVLSSMQHF SEAFHQVLGEKHKRGHLAEAEGHRDTCDEDS

研究背景	This gene encodes a C2H2 type zinc-finger protein by its similarity to the mouse Bcl11a/Evi9 protein. The corresponding mouse gene is a common site of retroviral integration in myeloid leukemia, and may functi on as a leukemia disease gene, in part, through its interaction with BCL6. During hematopoietic cell differ entiation, this gene is down-regulated. It is possibly involved in lymphoma pathogenesis since translocati ons associated with B-cell malignancies also deregulates its expression. Multiple transcript variants encod ing several different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
基因ID	53335
基因名	BCL11A
Swiss	Q9H165
别名	BCL11A-L; BCL11A-S; BCL11A-XL; BCL11a-M; CTIP1; DILOS; EVI9; HBFQTL5; ZNF856

产品验证



Western blot analysis of CTIP1 expressed in Rat brain using CTIP1 Rabbit mAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn