

Histone H2A Rabbit mAb

货号: B29396

产品信息

| 反应 | Human,Mouse,Rat |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 14kDa |
| 实测分子量 | 14kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HeLa,HepG2,Jurkat,Mouse spleen,Mouse thymus,Rat liver |
| 细胞定位 | Chromosome, Nucleus |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein corresponding to Human Histone H2A. |
|------|--|
| 序列 | MSGRGKQGGKARAKAKTRSSRAGLQFPVGRVHRLLRKGNYSERVGAGAPVYLAAVLEYLTAEILELAGNAARDNKKTRIIP RHLQLAIRNDEELNKLLGR |

靶点信息

| 研究背景 | Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octa mer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent his tone that is a member of the histone H2A family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015] |
|--------------|---|
| 基因 ID | 3012 |
| 基因名 | |
| Swiss | P04908 |
| 别名 | H2A.1; H2A.2; H2A/a; H2AFA |

产品验证



Western blot analysis of Histone H2A expressed in Mouse spleen,Rat liver,HeLa using Histone H2A Ra bbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure ti me: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn