

LOX Rabbit mAb

货号: B29452

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	46kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549
细胞定位	Secreted,extracellular space
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human LOX.
序列	MRFAWTVLLLGPLQLCALVHCAPPAAGQQQPPREPPAAPGAWRQQIQWENNGQVFSLLSLGSQYQPQRRRDPGAAVPG AANASAQQPRTPILLIRDNRTAAARTRTAGSSGVTAGRPRPTARHWFQAGYSTSRAREAGASRAENQTAPGEVPALSNLR PPSRVDGMVG


靶点信息

研究背景	This gene encodes a member of the lysyl oxidase family of proteins. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed to generate a regulatory propeptide and the mature enzyme. The copper-dependent amine oxidase activity of this enzyme functions in the crosslinking of collagens and elastin, while the propeptide may play a role in tumor suppression. In addition, defects in this gene have been linked with predisposition to thoracic aortic aneurysms and dissections.
基因ID	4015
基因名	LOX
Swiss	P28300
别名	LOX;AAT10

产品验证



Immunohistochemical analysis of paraffin-embedded human breast, using LOX Antibody.



Western blot analysis of LOX expressed in 293T,A-549 using LOX Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn