

TMEFF2 Rabbit mAb

货号: B29471

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	41kDa
实测分子量	41kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,PC-3,SH-SY5Y,Rat testis
细胞定位	basement membrane,extracellular region,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TMEFF2.
序列	HGKCEHSINMQEPSCRCDAGYTGQHCEKKDYSVLYVVPGPVRFQYVLIAAVIGTIQIAVICVVVLCITRKCPRSNRIHRQKQN TGHYSSDNTTRASTRLI

靶点信息

研究背景	This gene encodes a member of the tomoregulin family of transmembrane proteins. This protein has been shown to function as both an oncogene and a tumor suppressor depending on the cellular context and m ay regulate prostate cancer cell invasion. Multiple soluble forms of this protein have been identified that arise from both an alternative splice variant and ectodomain shedding. Additionally, this gene has been f ound to be hypermethylated in multiple cancer types. Alternative splicing results in multiple transcript va riants. [provided by RefSeq, Feb 2015]
------	---

基因ID	23671
基因名	TMEFF2
Swiss	Q9UIK5
别名	CT120.2; HPP1; TENB2; TPEF; TR; TR-2

产品验证



Western blot analysis of TMEFF2 expressed in HeLa,PC-3,SH-SY5Y using TMEFF2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Block ing buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn