

DOCK2 Rabbit mAb

货号: B29518

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	38kDa/211kDa
实测分子量	212kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat
细胞定位	Cytoplasm,Endomembrane system,Peripheral membrane protein,cytoskeleton
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human DOCK2.
序列	FTEEYVRDHPEDQDKLTHLKDLIAWQIPFLGAGIKIHEKRVSDNLRPFHDRMEECFKNLKMKVEKEYGVREMPDFDDRRVGRPRSMRLRSYRQMSIISLASMNSDCSTPSKPTSESFDELELASPKTPRVEQEEPISPGSTLPEVKLRRSKKRTKRSSVFADEKAAAESDLKRLSRKHEFMSDTNLSEHAAIPLKASVLSQMSFASQSMPTIPALALSVAGIPGLDEANTSPRLSQTFLQLSDGDKKTLTRKKVNQFFKTMLASKSAEEGKQIPDSLSTD

靶点信息

研究背景	The protein encoded by this gene belongs to the CDM protein family. It is specifically expressed in hematopoietic cells and is predominantly expressed in peripheral blood leukocytes. The protein is involved in remodeling of the actin cytoskeleton required for lymphocyte migration in response to chemokine signaling . It activates members of the Rho family of GTPases, for example RAC1 and RAC2, by acting as a guanine nucleotide exchange factor (GEF) to exchange bound GDP for free GTP.
------	---

基因ID	1794
基因名	DOCK2
Swiss	Q92608
别名	DOCK2;IMD40

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn