

Mad2L1 Rabbit mAb

货号: B29577

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	10kDa/23kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver,Rat liver
细胞定位	Chromosome,Cytoplasm,Nucleus,centromere,cytoskeleton,kinetochore,spindle pole
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Mad2L1.
序列	MALQLSREQGITLRGSAEIVAEFFSFGINSILYQRGIYPSETFTRVQKYGLTLLVTTDLELIKYLNNVVEQLKDWLYKCSVQKL VVVISNIESGEVLERWQFDIECDKTAKDDSAPREKSQKAIQDEIRSVIRQITATVTFLPLEVSCSFDLLIYTDKDLVVPEKWE ESGPQFITNSEEVRLRSFTTTIHKVNSMVAYKIPVND

靶点信息

研究背景	MAD2L1 is a component of the mitotic spindle assembly checkpoint that prevents the onset of anaphase until all chromosomes are properly aligned at the metaphase plate. MAD2L1 is related to the MAD2L2 gene located on chromosome 1. A MAD2 pseudogene has been mapped to chromosome 14.
基因ID	4085

基因名	MAD2L1
Swiss	Q13257
别名	MAD2L1;HSMAD2;MAD2

产品验证



Western blot analysis of Mad2L1 expressed in Mouse liver,Rat liver using Mad2L1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn