

Eph receptor B3 Rabbit mAb

货号: B29622

产品信息

uman,Mouse,Rat
abbit
onoclonal
'B IF/ICC IP FC
B: 1:500 - 1:2000 /ICC: 1:50 - 1:200 : 1:20 - 1:50 C: 1:20 - 1:50
.0kDa
.0kDa
quid
ore at -20°C. Avoid freeze / thaw cycles. Iffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
nconjugated
eLa,HT-29,SH-SY5Y,Mouse brain,Rat brain
ell membrane,Cell projection,Single-pass type I membrane protein,dendrite
finity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Eph receptor B3.
序列	SELAWTSHPESGWEEVSGYDEAMNPIRTYQVCNVRESSQNNWLRTGFIWRRDVQRVYVELKFTVRDCNSIPNIPGSCKET FNLFYYEADSDVASASSPFWM

靶点信息

研究背景	Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, an d the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided i nto two groups based on the similarity of their extracellular domain sequences and their affinities for bind ing ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. This gene encodes a receptor for ephrin-B family members.
基因 ID	2049
基因名	ЕРНВЗ
Swiss	P54753
别名	EPHB3; EK2; ETK2; HEK2; TYRO6; EPH receptor B3

产品验证



Western blot analysis of Eph receptor B3 expressed in Mouse brain,Rat brain,HeLa using Eph recept or B3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysate s/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit . Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn