

FADS1 Rabbit mAb

货号: B29649

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	42kDa/51kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse testis,Rat lung
细胞定位	mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human FADS1.
序列	YNISEFTRRHPGGSRVISHYAGQDATDPFVAFHINKGLVKKYMNSLLIGELSPEQPSFEPTKNKELTDEFRELRATVERMGL MKANHVFFLLYLLHILLLD

靶点信息

研究背景	The protein encoded by this gene is a member of the fatty acid desaturase (FADS) gene family. Desaturas e enzymes regulate unsaturation of fatty acids through the introduction of double bonds between defined carbons of the fatty acyl chain. FADS family members are considered fusion products composed of an N-t erminal cytochrome b5-like domain and a C-terminal multiple membrane-spanning desaturase portion, b oth of which are characterized by conserved histidine motifs. This gene is clustered with family members FADS1 and FADS2 at 11q12-q13.1; this cluster is thought to have arisen evolutionarily from gene duplicat ion based on its similar exon/intron organization.
基因ID	3992
基因名	FADS1
Swiss	O60427
别名	FADS1;D5D;FADS6;FADSD5;LLCDL1;TU12

产品验证



Western blot analysis of FADS1 expressed in 293T, Mouse testis, Rat lung using FADS1 Rabbit mAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn