

MEK2 Rabbit mAb

货号: B29672

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	44kDa
实测分子量	44kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,HT-29
细胞定位	Cytoplasm,Membrane,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human MEK2.
序列	MLARRKPVLPAITNPTIAEGPSPTSEGASEANLVDLQKKLEELELDEQKKRLEAFLTQKAKVGELKDDDFERISELGAGN GGVTKVQHRPSGLIMAR

靶点信息

研究背景	The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is known to play a critical role in mitogen growth factor signal transduction. It phosphorylates and thus activates MAPK1/ERK2 and MAPK2/ERK3. The activation of this kinase itself is dependent on the Ser/Thr phosphorylation by MAP kinase kinase kinases. Mutations in this gene cause cardiofacio cutaneous syndrome (CFC syndrome), a disease characterized by heart defects, mental retardation, and distinctive facial features similar to those found in Noonan syndrome. The inhibition or degradation of this kinase is also found to be involved in the pathogenesis of Yersinia and anthrax. A pseudogene, which is located on chromosome 7, has been identified for this gene.
基因ID	5605
基因名	MAP2K2
Swiss	P36507
别名	CFC4;MAPKK2;MEK2;MKK2;PRKMK2;MAP2K2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn