

Nodal Rabbit mAb

货号: B29698

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	39kDa
实测分子量	43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,HepG2,293T,Rat liver
细胞定位	Secreted
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Nodal.
序列	SNLSQEQRQLGGSTLLWEAESSWRAQEGQLSWEWGKRHRRHHLPDRSQLCRKVKFQVDFNLIGWGSWIIYPKQYNAY RCEGECPNPVGEEFHPTNHAYIQSLLKRYQPHRVPSTCCAPVKTKPLSMLYVDNGRVLLDHHKDMIVEECGCL


靶点信息

研	究	背	景

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of pro teins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMA D family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate the mature protein, which regulates early embryonic development. This protein is required for maintenance of human embryonic stem cell pluripotency and may play a role in human place ntal development. Mutations in this gene are associated with heterotaxy, a condition characterized by ran dom orientation of visceral organs with respect to the left-right axis.

基因ID	4838
基因名	NODAL
Swiss	Q96S42
别名	NODAL;HTX5

产品验证



Western blot analysis of Nodal expressed in SH-SY5Y,HepG2,293T using Nodal Rabbit mAb at 1:1000. S econdary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blockin g buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn