

Phospho-Stat5 (Y694) Rabbit mAb

货号: B29708

产品信息

| 反应 | Human,Mouse |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 |
| 理论分子量 | 90kDa |
| 实测分子量 | 90kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | TF-1 |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein corresponding to Human Phospho-Stat5 (Y694). |
|------|---|
| 序列 | DGYVK |

靶点信息

| 研究背景 | The protein encoded by this gene is a member of the STAT family of transcription factors. In response to c ytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinase s, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcripti on activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, I L3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and ha s been shown to be essential for tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2013] |
|--------------|---|
| 基因 ID | 6776 |
| 基因名 | STAT5A |
| Swiss | P42229 |
| 别名 | MGF;STAT5;STAT5A |

产品验证



Western blot analysis of Phospho-Stat5 (Y694) expressed in TF-1 using Phospho-Stat5 (Y694) Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 12 0s.

实验步骤

访问官网浏览详情: www.ablybio.cn