

Cdk4 Rabbit mAb

货号: B29721

产品信息

| 反应 | Human |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC FC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50 |
| 理论分子量 | 20kDa/33kDa |
| 实测分子量 | 34kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T,HeLa |
| 细胞定位 | Cytoplasm,Membrane,Nucleus |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein corresponding to Human Cdk4. |
|------|--|
| 序列 | VGCIFAEMFRRKPLFCGNSEADQLGKIFDLIGLPPEDDWPRDVSLPRGAFPPRGPRPVQSVVPEMEESGAQLLLEMLTFNP HKRISAFRALQHSYLHKDEGNPE |

靶点信息

| 研究背景 | The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of S. cerevisiae cdc28 and S. pombe cdc2. It is a catalytic subunit of the prot ein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is rest ricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p 16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene product (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and R b were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation site s of this gene have been reported. |
|-------|--|
| 基因ID | 1019 |
| 基因名 | CDK4 |
| Swiss | P11802 |
| 别名 | CDK4;CMM3;PSK-J3 |

产品验证



Western blot analysis of Cdk4 expressed in 293T,HeLa using Cdk4 Rabbit mAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn