

FEN1 Rabbit mAb

货号**: B29763**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	42kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HUVEC,Mouse thymus
细胞定位	Mitochondrion, Nucleus, nucleoplasm
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human FEN1.	
序列	IQKHKSIEEIVRRLDPNKYPVPENWLHKEAHQLFLEPEVLDPESVELKWSEPNEEELIKFMCGEKQFSEERIRSGVKRLSKSR QGSTQGRLDDFFKVTGSL	

靶点信息

研究背景	The protein encoded by this gene removes 5' overhanging flaps in DNA repair and processes the 5' ends of Okazaki fragments in lagging strand DNA synthesis. Direct physical interaction between this protein an d AP endonuclease 1 during long-patch base excision repair provides coordinated loading of the proteins onto the substrate, thus passing the substrate from one enzyme to another. The protein is a member of t he XPG/RAD2 endonuclease family and is one of ten proteins essential for cell-free DNA replication. DNA secondary structure can inhibit flap processing at certain trinucleotide repeats in a length-dependent ma nner by concealing the 5' end of the flap that is necessary for both binding and cleavage by the protein e ncoded by this gene. Therefore, secondary structure can deter the protective function of this protein, lea ding to site-specific trinucleotide expansions. [provided by RefSeq, Jul 2008]
基因ID	2237
基因名	FEN1
Swiss	P39748
别名	FEN-1; MF1; RAD2

产品验证



Western blot analysis of FEN1 expressed in 293T,HUVEC,Mouse thymus using FEN1 Rabbit mAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bloc king buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn