

# **Actin Rabbit mAb**

货号: B29771

## 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	42kDa
实测分子量	42kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,A-431,RD,C6,Mouse lung,Mouse brain,Mouse heart,Rat lung,Rat heart
细胞定位	Cytoplasm,cytoskeleton
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Actin.
序列	MCDEDETTALVCDNGSGLVKAGFAGDDAPRAVFPSIVGRPRHQGVMVGMGQKDSYVGDEAQSKRGILTLKYPIEHGIIT NWDDMEKIWHHTFYNELRVAP

## 靶点信息

研究背景	The product encoded by this gene belongs to the actin family of proteins, which are highly conserved prot eins that play a role in cell motility, structure and integrity. Alpha, beta and gamma actin isoforms have b een identified, with alpha actins being a major constituent of the contractile apparatus, while beta and gamma actins are involved in the regulation of cell motility. This actin is an alpha actin that is found in skele tal muscle. Mutations in this gene cause nemaline myopathy type 3, congenital myopathy with excess of thin myofilaments, congenital myopathy with cores, and congenital myopathy with fiber-type disproportion, diseases that lead to muscle fiber defects. [provided by RefSeq, Jul 2008]
------	--

基因ID	58
基因名	ACTA1
Swiss	P68133
别名	ACTA; ASMA; CFTD; CFTD1; CFTDM; MPFD; NEM1; NEM2; NEM3; SHPM

## 产品验证



Western blot analysis of Actin expressed in Mouse lung,Rat lung,HeLa using Actin Rabbit mAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Block ing buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn