

# Zic1 Rabbit mAb

货号: B29784

## 产品信息

| 反应    | Human,Mouse,Rat  |
|-------|--|
| 宿主    | Rabbit   |
| 克隆性   | Monoclonal   |
| 预测反应  |  |
| 应用    | WB IF/ICC FC   |
| 推荐浓度  | WB: 1:500 - 1:2000<br>IF/ICC: 1:50 - 1:200<br>FC: 1:20 - 1:50                              |
| 理论分子量 |  |
| 实测分子量 | 48kDa  |
| 形式    | Liquid   |
| 保存条件  | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物   | Unconjugated   |
| 阳性对照  | U-87MG,U-251MG,Mouse brain,Rat brain   |
| 细胞定位  | cytoplasm,nucleoplasm,nucleus  |
| 纯化    | Affinity purification  |

## 抗原信息


| 抗原信息 | Recombinant fusion protein corresponding to Human Zic1.   |
|------|---|
| 序列   | LFAASAGGFGGPHGHTDAAGHLLFPGLHEQAAGHASPNVVNGQMRLGFSGDMYPRPEQYGQVTSPRSEHYAAPQLHG<br>YGPMNVNMAAHHGA |

## 靶点信息

| 研究背景 | This gene encodes a member of the ZIC family of C2H2-type zinc finger proteins. Members of this family are important during development. Aberrant expression of this gene is seen in medulloblastoma, a childh ood brain tumor. This gene is closely linked to the gene encoding zinc finger protein of the cerebellum 4, a related family member on chromosome 3. This gene encodes a transcription factor that can bind and tr ansactivate the apolipoprotein E gene. [provided by RefSeq, Jul 2008] |
|------|--|
|------|--|

| 基因ID  | 7545                 |
|-------|----------------------|
| 基因名   | ZIC1                 |
| Swiss | Q15915               |
| 别名    | CRS6;ZIC;ZNF201;ZIC1 |

## 产品验证



Western blot analysis of Zic1 expressed in Mouse brain,Rat brain,U-87MG using Zic1 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn