

NCAM Rabbit mAb

货号: B29814

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	40kDa/73kDa/80kDa/83kDa/93kDa/94kDa
实测分子量	180kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Mouse brain,Rat brain
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NCAM.
序列	SEEKASWTRPEKQETLDGHMVVRSHARVSSLTKSIQYTDAGEYICTASNTIGQDSQSMYLEVQYAPKLQGPVAVYTWEG NQVNITCEVFAYPSATISWFRDGQLLPSSNYSNIKIYNTPSASYLEVTPDSENDGFGNYNCTAVNRIGQESLEFILVQADTPSS PSIDQVEPYSSTAQVQFDEPEATGGVPILKYKAEWRAVGEEVWHSKWYDAKEASMEGIVTIVGLKPETTYAVRLAALNGK GLGEISAASEFKTQPVQGEPSAPKLEGQMGEDGNS

靶点信息

研究背景	This gene encodes a cell adhesion protein which is a member of the immunoglobulin superfamily. The encoded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein has been shown to be involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance. Alternative splicing results in multiple transcript variants.
基因ID	4684
基因名	NCAM1
Swiss	P13591
别名	CD56;MSK39;NCAM;NCAM1;NCAM1 /CD56

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn