

IKK gamma Rabbit mAb

货号: B29940

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	36kDa/48kDa/55kDa
实测分子量	48kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,Mouse liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human IKK gamma.
序列	MNRHLWKSQLCCEMVQPSGGPAADQDVLGEESPLGKPAMLHLPSEQGAPETLQRCLEENQELRDAIRQSNQILRERCEELL HFQASQREEKEFLMCKFQEARKLVERLGLEKLDLKRQKEQALREVEHLKRCQQQMAEDKASVKAQVTSLLGELQESQSRL EAATKECQALEGRARAASEQARQLESEREALQQQHSVQVDQLRMQQQSVEAALRMERQAASEEKRKLAQLQVAYHQLF QEYDNHIKSSVVGSEKRGMQLEDLKQQLQQAEEALVAKQEVIDKLKEEAQHKIVMETVPVLKAQADIYKADFQAERQA REKLAIEKKELLQEQLQREYSKLKASCQESARIEDMRKRHVEVSQAPLPAPAYLSSPLALPSQRRSPPEEPPDFCCPKC QYQAPDMDTLQIHVMECIE

靶点信息

研究背景	This gene encodes the regulatory subunit of the inhibitor of kappaB kinase (IKK) complex, which activates NF-kappaB resulting in activation of genes involved in inflammation, immunity, cell survival, and other pathways. Mutations in this gene result in incontinentia pigmenti, hypohidrotic ectodermal dysplasia, and several other types of immunodeficiencies. A pseudogene highly similar to this locus is located in an adjacent region of the X chromosome.
基因ID	8517
基因名	IKBKG
Swiss	Q9Y6K9
别名	IKBKG;AMCBX1;FIP-3;FIP3;Fip3p;IKK-gamma;IKKAP1;IKKG;IMD33;IP;IP1;IP2;IPD2;NEMO;ZC2HC9

产品验证



Western blot analysis of IKK gamma expressed in A-549,Mouse liver using IKK gamma Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn