

PD-L1 (CD274) Rabbit mAb

货号: B29957

产品信息

反应	Human, Mouse, Rat
2.0-	Trainally lease, lac
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	20kDa/33kDa
实测分子量	46kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2
细胞定位	Cell membrane,Endomembrane system,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PD-L1 (CD274).
序列	NAPYNKINQRILVVDPVTSEHELTCQAEGYPKAEVIWTSSDHQVLSGKTTTTNSKREEKLFNVTSTLRINTTTNEIFYCTFRRL DPEENHTAELVIPELPLAHPPNERT

靶点信息

研究背景	This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hem atopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand wit h its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of norma I tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the imm une response. In tumor microenvironments, this interaction provides an immune escape for tumor cells th rough cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in multiple transcript variants.
基因 ID	29126
基因名	CD274
Swiss	Q9NZQ7
别名	CD274; B7-H; B7H1; PD-L1; PDCD1L1; PDCD1LG1; PDL1; CD274 molecule

产品验证



Western blot analysis of PD-L1 (CD274) expressed in HepG2 using PD-L1 (CD274) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blockin g buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn