

Fibronectin Rabbit mAb

货号: B30000

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	71-73kDa/221-272kDa
实测分子量	263kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,NIH/3T3
细胞定位	Secreted,extracellular matrix,extracellular space
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Fibronectin.
序列	LLCQCLGFSGHFRCDSSRWCHDNGVNYKIGEKWDRQGENGQMMSTCLGNKGKFCDPHEATCYDDGKTYHVGE QWQKEYLGAICSCTCFGQRGWRCNDNCRRPGGEPSPEGTTGQSNTQYSQRYHQRTNTNVNCPIECFMPLDVQADRED SRE

靶点信息

研究背景	This gene encodes fibronectin, a glycoprotein present in a soluble dimeric form in plasma, and in a dimeric or multimeric form at the cell surface and in extracellular matrix. The encoded preprotein is proteolytically processed to generate the mature protein. Fibronectin is involved in cell adhesion and migration processes including embryogenesis, wound healing, blood coagulation, host defense, and metastasis. The gene has three regions subject to alternative splicing, with the potential to produce 20 different transcript variants, at least one of which encodes an isoform that undergoes proteolytic processing. The full-length nature of some variants has not been determined.
基因ID	2335
基因名	FN1
Swiss	P02751
别名	FN1;CIG;ED-B;FINC;FN;FNZ;GFND;GFND2;LETS;MSF;fibronectin

产品验证



Western blot analysis of Fibronectin expressed in HepG2, NIH/3T3 using Fibronectin Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn