

TNF beta Rabbit mAb

货号: B30015

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	22kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse spleen
细胞定位	Membrane,Secreted
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TNF beta.
序列	LPGVGLTPSAAQTARQHPKMHLAHSTLKPAAHLIGDPSKQNSLLWRANTDRAFLQDGFSLSNNSLLVPTSGIYFVYSQVV FSGKAYSPKATSSPLYLAHEVQLFSSQYPFHVPLLSSQKMVYPGLQEPWLHSMYHGAAFQLTQGDQLSTHTDGIPHLVLS PSTVFFGAFAL

靶点信息

	The encoded protein, a member of the tumor necrosis factor family, is a cytokine produced by lymphocyt
	es. The protein is highly inducible, secreted, and forms heterotrimers with lymphotoxin-beta which ancho
	r lymphotoxin-alpha to the cell surface. This protein also mediates a large variety of inflammatory, immu
	nostimulatory, and antiviral responses, is involved in the formation of secondary lymphoid organs during
	development and plays a role in apoptosis. Genetic variations in this gene are associated with susceptibili
	ty to leprosy type 4, myocardial infarction, non-Hodgkin's lymphoma, and psoriatic arthritis. Alternatively
	spliced transcript variants have been observed for this gene.

基因 ID	4049
基因名	LTA
Swiss	P01374
别名	LTA;LT;TNFB;TNFSF1;TNLG1E;LT-alpha

产品验证



Western blot analysis of TNF beta expressed in Mouse spleen using TNF beta Rabbit mAb at 1:1000. Seco ndary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buff er: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn