

Rhodopsin Rabbit mAb

货号: B30035

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	39kDa
实测分子量	39kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse eye,Rat eye
细胞定位	Membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Rhodopsin.
序列	EVTRMVIIMVIAFLICWVPYASVAFYIFTHQGSNFGPIFMTIPAFFAKSAAIYNPVIYIMMNKQFRNCMLTTICCGKNPLGDDE ASATVSKTETSQVAPA

靶点信息

研究背景	Retinitis pigmentosa is an inherited progressive disease which is a major cause of blindness in western co mmunities. It can be inherited as an autosomal dominant, autosomal recessive, or X-linked recessive diso rder. In the autosomal dominant form, which comprises about 25% of total cases, approximately 30% of fa milies have mutations in the gene encoding the rod photoreceptor-specific protein rhodopsin. This is the t ransmembrane protein which, when photoexcited, initiates the visual transduction cascade. Defects in thi s gene are also one of the causes of congenital stationary night blindness. [provided by RefSeq, Jul 2008]
------	--

基因ID	6010
基因名	RHO
Swiss	P08100
别名	CSNBAD1; OPN2; RP4

产品验证



Western blot analysis of Rhodopsin expressed in Mouse eye,Rat eye using Rhodopsin Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn