

# Phospho-SIRT1 (S47) Rabbit mAb

货号: B30039

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200
理论分子量	61kDa/81kDa
实测分子量	82kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Mitochondrion,Nucleus,Nucleus,PML body
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Phospho-SIRT1 (S47).
序列	TILKDLLPETIPPPELDDMTLWQIVINILSEPPKRKKRKDINTIEDAVKLLQECKKIIVLTGAGVSVSCGIPDFRSRDGIYARLAV DFPDLPDPQAMFDIE

靶点信息

研究背景	This gene encodes a member of the sirtuin family of proteins, homologs to the yeast Sir2 protein. Membe rs of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. The funct ions of human sirtuins have not yet been determined; however, yeast sirtuin proteins are known to regul ate epigenetic gene silencing and suppress recombination of rDNA. Studies suggest that the human sirtuins may function as intracellular regulatory proteins with mono-ADP-ribosyltransferase activity. The protein encoded by this gene is included in class I of the sirtuin family. Alternative splicing results in multiple transcript variants.
基因ID	23411
基因名	SIRT1
Swiss	Q96EB6
别名	SIR2;SIR2L1;SIR2alpha;SIRT1;sirtuin 1

#### 产品验证



Western blot analysis of Phospho-SIRT1 (S47) expressed in 293T using Phospho-SIRT1 (S47) Rabbit mA b at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 c

#### 实验步骤

访问官网浏览详情: www.ablybio.cn